

## Bibliography of documents about Seagrass Communities in Florida

Associates, C. S. (1992). Roller-frame trawl bait shrimp fishery impacts on by-catch, seagrass meadows, and habitat quality in Pine Island Sound, Florida, Florida Marine Research Institute: 42.

Avery, W. and R. Johansson (2002). Data summary from the Tampa Bay Interagency Seagrass Monitoring Program through the year 2001. Tampa, FL, Tampa Bay Estuary Program: 73.

Brook, I. M. (1978). "Comparitive macrofaunal abundance in turtlegrass (*Thalassia testudinum*) communities in South Florida characterized by blade density." *Bulletin of Marine Science* 28(1): 212-217.

Camp, D. K., S. B. Cobb, et al. (1973). "Overgrazing of seagrasses by a regular urchin, *Lytechinus variegatus*." *Florida Department of Natural Resources Marine Research Laboratory* 23(1): 37-38.

Cross, L. (2007). Feather Sound seagrass recovery project: Final report and management recommendations. St. Petersburg, FL: Tampa Bay Estuary Program. Technical Report 03-07.

Darovec, J. E., Jr., Carlton, J. M., et al. (1975). "Techniques for coastal restoration and fishery enhancement in Florida." St. Petersburg, FL, Florida Department of Natural Resources, Marine Research Laboratory. Florida Marine Research Publications, 15.

Dawes, C. J., Hall, M. O., et al. (1985). "Seasonal biomass and energy content in seagrass communities on the West Coast of Florida." *Journal of Coastal Research*, 1(3): 255-262.

Dawes, C. J., Phillips, R. C., et al. (2004). Seagrass communities of the Gulf Coast of Florida: status and ecology. St. Petersburg, FL, Florida Fish and Wildlife Conservation Commission Fish and Wildlife Research Institute and the Tampa Bay Estuary Program.

Dixon, L. K., and Kirkpatrick, G. J. (1995). Light attenuation with respect to seagrasses in Sarasota Bay. Sarasota, FL: Mote Marine Laboratory. Technical Report.

Dixon, L. K., and Kirkpatrick, G. J. (1999). Causes of light attenuation with respect to seagrasses in upper and lower Charlotte Harbor. Mote Marine Laboratory Technical Report to the Charlotte Harbor National Estuary Program and the Southwest Florida Water Management District.

Dixon, L. K., and Leverone, J. (1995). Light requirements of *Thalassia* in Tampa Bay. Sarasota, FL: Mote Marine Laboratory. Technical Report.

Dixon, L. K., and Perry, J. S. (2003). Analysis of changes in seagrass coverage (1996-2001). Sarasota, FL: Mote Marine Laboratory. Technical Report 650.

Dixon, L. K., Hall, and Kirkpatrick, G. J. (2010). A spectrally explicit optical model of attenuation for Charlotte Harbor seagrasses. Sarasota, FL: Mote Marine Laboratory. Technical Report 1460.

Dixon, L. K. and Kirkpatrick, G. J. (1999). Causes of light attenuation with respect to seagrasses in Upper and Lower Charlotte Harbor. Sarasota, FL: Mote Marine Laboratory.

Dixon, L. K. and Kirkpatrick, G. J. (1995). Light attenuation with respect to seagrasses in Sarasota Bay, Florida. Sarasota, FL: Mote Marine Laboratory.

Dixon, L. K. and Perry, J. S. (2003). Analysis of Changes in Seagrass Coverage, 1996-1999 and 1999-2001, for Anna Maria Sound, Sarasota, Roberts, Little Sarasota, and Blackburn Bays. Sarasota, FL: Mote Marine Laboratory: 38 plus Appendices.

Durako, M. J. (1993). New approaches to seagrass restoration. St. Petersburg, FL: Florida Marine Research Institute.

Folit, R. and Morris, J. (1992). Beds, Boats and Buoys: a Study in Protecting Seagrass Beds from Motorboat Propeller Damage. Sarasota, FL, New College of University of Florida: 45 plus Appendices.

Fonseca, M. S. (1994). A guide to planting seagrasses in the Gulf of Mexico, Texas A&M Sea Grant Institution: 25.

Fonseca, M. S., Robbins, B. D., et al. Evaluating the effect of offshore sandbars on seagrass recovery and restoration in Tampa Bay through ecological forecasting and hindcasting of exposure to waves, NOAA Center for Coastal Fisheries and Habitat Research: 49.

Fourqurean, J. W. and Zieman, J. C. (1991). "Photosynthesis, respiration, and whole plant carbon budget of the seagrass *Thalassia testudinum*." Marine Ecology Progress Series 69: 161-170.

Gilbert, S., Robbins, B.D., Luther, M., and Lewis, R. (2009). Field measurements of bottom stress due to wave action and tidal velocity in Tampa Bay to support seagrass restoration and shoreline protection. TBEP Tech Publ. 03-09.

Gilmore, R. G., Donohoe, C. J., Cooke, D. W. (1983). "Observations on the distribution and biology of east-central florida populations of the common snook, *Centropomus undecimalis* (bloch)." Academy Symposium 46(3/4): 306-313.

Greening, H. (1995). Tampa Bay Atmospheric Deposition Study Great Waters Program, Tampa Bay National Estuary Program.

Greening, H., editor. (2000). *Seagrass Management: It's Not Just Nutrients! Proceedings of a Symposium*. St. Petersburg, FL: Tampa Bay Estuary Program.

Greening, H. (2004). Factors influencing seagrass recovery in Feather Sound, Tampa Bay, Florida, Pinellas County Environmental Foundation: 29.

Greening, H. (2006). Toward reversal of eutrophic conditions in a subtropical estuary: water quality and seagrass response to nitrogen loading reductions in Tampa Bay, Florida, USA. Environmental Management.

Heck, K. L., Jr. (1979). "Some determinants of the composition and abundance of motile macroinvertebrates species in tropical and temperate turtlegrass (*Thalassia testudinum*) meadows." Journal of Biogeography 6: 183-200.

Heck, K. L., Jr., and Wetstone, G. S. (1977). "Habitat complexity and invertebrate species richness and abundance in tropical seagrass meadows." *Journal of Biogeography* 4: 135-142.

Heffernan, J. J., Gibson, R.A. (1983). "A comparison of primary production rates in Indian river, Florida seagrass systems." *Florida Scientist* 46(3/4): 286-295.

Kurz, R. C., Tomasko, D., et al. (1999). Summary of recent trends in seagrass distributions in southwest Florida coastal waters. Tampa, FL, SWIM/SFWMD.

Kurz, R. C., Fenwick, R. W., et al. (1998). A comparison of fish communities in restored and natural salt marshes in Tampa Bay, Florida. Tampa, Florida, SWIM Section, SFWMD.

Leber, K. M. (1985). "The influence of predatory decapods, refuge, and microhabitat selection on seagrass communities." *Ecology* 66(6).

Leber, K. M., and Greening, H. (1985). "Community studies in seagrass meadows: A comparison of two methods for sampling macroinvertebrates and fishes." *Fishery Bulletin* 84(2).

Leverone, J. R. and Marshall, M. J. (1992). Seagrass Bed Faunal Utilization Study. Sarasota, FL, Mote Marine Laboratory.

Lewis, F. G., III (1984). "Distribution of macrobenthic crusteans associated with *Thalassia*, *Halodule* and bare sand substrata." *Marine Ecology Progress Series* 19: 101-113.

Livingston, R. J. (1984). "The relationship of physical factors and biological response in coastal seagrass meadows." *Estuaries* 7(4A): 377-390.

McMillan, C. (1980). "Reproductive physiology in the seagrass, *Syringodium filiforme*, from the Gulf of Mexico and the Caribbean." *American Journal of Botany* 67(1): 104-110.

McMillan, C., and Phillips, R. C. (1979). "Differentiation in habitat response among populations of New World seagrasses." *Aquatic Botany* 7: 185-196.

Meyer, D. L., et al. (1999). The impact of bait shrimp trawling on seagrass beds and fish by-catch in Tampa Bay, Florida. *Fishery Bulletin*: 97(1): 193-199.

Milbrandt, E. C., (2009). The effects of root zone manipulation on microcosm *Thalassia testudinum* transplants. *Florida Scientist* 72:406-417.

Milbrandt, E. C., Greenawalt, M, Sokoloff, P. O. (2008). Short-term indicators of seagrass transplant stress in response to rhizosphere and bacterial community disruption. *Botanica Marina* 51:103-111.

Morgan, M. D. and Kitting, C. L. (1984). "Productivity and utilization of the seagrass *Halodule wrightii* and its attached epiphytes." *Limnology and Oceanography* 29(5): 1066-1076.

Orth, R. J., Heck, K. L. Jr., et al. (1984). "Faunal communities in seagrass beds: a review of the influence of plant structure and prey characteristics on predator-prey relationships." *Estuaries* 7: 339-350.

Pangallo, R. A., Bell, S.S. (1988). "Dynamics of the aboveground and belowground structure of the seagrass *Halodule wrightii*." Marine Biology Progress Series 43: 297-301.

Patriquin, D. G. (1975). "'Migration' of blowouts in seagrass beds at Barbados and Carriacou, West Indies, and its ecological and geological implications." Aquatic Botany 1: 163-189.

Phillips, R. C. (1960). "Environmental effect on leaves of *Diplanthera du Petit-Thouars*." Bulletin of Marine Science of the Gulf and Caribbean 10(3): 346-353.

Rice, J. D., and Trocine, R. P. (1983). "Factors influencing seagrass ecology in the Indian River Lagoon." Florida Scientist 46(3/4): 250-276.

Robbins, B. D. (1997). "Quantifying temporal change in seagrass areal coverage: the use of GIS and low resolution aerial photography." Aquatic Botany 58: 259-267.

Robbins, B. D., and Bell, S. S. (2000). "Dynamics of a subtidal seagrass landscape: seasonal and annual change in relation to water depth." Ecology 81: 1193-1205.

Ryer, C. H., Montfrans, J. V. (1990). "Utilization of a seagrass meadow and tidal marsh creek by blue crabs *Callinectes sapidus*. II. Spatial and temporal patterns of molting." Bulletin of Marine Science 46(1): 95-104.

Sargent, F. J., Leary, T. J., et al. (1995). Scarring of Florida's seagrasses: assessment and management options. St. Petersburg, FL, Florida Department of Environmental Protection Florida Marine Research Institute: 43 plus appendices.

Serviss, G. M. and Sauers, S. (2003). Sarasota Bay Juvenile Fisheries Habitat Assessment, SBEP: 212.

Sheng, Y. P., and Yassuda, E. A. (1995). Modeling the impact of nutrient load reduction on water quality and seagrass in Roberts Bay and Little Sarasota Bay.

Stoner, A. W. (1980). "Perception and choice of substratum by epifaunal amphipods associated with seagrasses." Marine Ecology Progress Series 3: 105-111.

Stoner, A. W. (1980). "The role of seagrass biomass in the organization of benthic macrofaunal assemblages." Bulletin of Marine Science 30(3): 537-551.

Stoner, A. W. (1983). "Distribution of fishes in seagrass meadows: role of macrophyte biomass and species composition." Fishery Bulletin 81(4): 837-846.

Southwest Florida Water Management District. (1992). Tampa Bay Surface Water Improvement and Management Plan. Brooksville, FL: Southwest Florida Water Management District.

Southwest Florida Water Management District. (1997). Sarasota Bay Surface Water Improvement and Management (SWIM) Plan. Brooksville, FL: Southwest Florida Water Management District.

- Taylor, J. L., and Saloman, C. H. (1968). "Some effects of hydraulic dredging and coastal development in Boca Ciega Bay, Florida." *Fishery Bulletin* 67(2): 213-241.
- Thorhaug, A. (1985). "Large-scale seagrass restoration in a damaged estuary." *Marine Pollution Bulletin* 16(2): 55-62.
- Thorne, R. F. (1995). "Vascular plants of Fort DeSoto Park, Pinellas County, Florida." *Selbyana* 16(1): 100-109.
- Tomasko, D. A., and Dawes, C. J. (1996). The effects of anthropogenic nutrient enrichment on turtle grass (*Thalassia testudinum*) in Sarasota Bay, Florida. *Estuaries and Coasts*.
- Tomasko, D. A., Corbett, C. A., Greening, H. S., and Raulerson, G. E. (2005). Spatial and temporal variation in seagrass coverage in Southwest Florida: assessing the relative effects of anthropogenic nutrient load reductions and rainfall in four contiguous estuaries. *Mar. Poll. Bull.*
- Tomasko, D. A. (1996). Report on status and trends of seagrass coverage in Sarasota Bay - 1988 to 1995.
- Tomasko, D. A., Lapointe, Brian E. (1991). "Productivity and biomass of *Thalassia testudinum* as related to water column nutrient availability and epiphyte levels: field observations and experimental studies." *Marine Ecology Progress Series* 75: 9-17.
- Tomasko, D. A., and Ries, T. F. (unknown) Responses of Tampa Bay and Sarasota Bay seagrass meadows to nitrogen load reductions.
- Tomasko, D., C. A. Corbett, et al. (2005). "Spatial and temporal variation in seagrass coverage in Southwest Florida: assessing the relative effects of anthropogenic nutrient load reductions and rainfall in four contiguous estuaries." *Marine Pollution Bulletin* 50: 797-805.
- Tomasko, D., Bristol, D. L., et al. (2001). "Assessment of present and future nitrogen loads, water quality, and seagrass (*Thalassia testudinum*) depth distribution in Lemon Bay, Florida." *Estuaries* 24(6A): 926-938.
- Tomlinson, P. B., Vargo, G.A. (1966). "On the morphology and anatomy of turtle grass, *Thalassia testudinum* (hydrocharitacean) I. Vegetative morphology." *Bulletin of Marine Science* 16(4): 749-761.
- Virnstein, R. W., and Mikkelsen, P. S. (1983). "Seagrass beds versus sand bottoms: The trophic importance of their associated benthic invertebrates." *Florida Science* 46(3/4): 355-362.
- Weinstein, M. P., and Heck, K.L., Jr. (1979). "Ichthyofauna of seagrass meadows along the Caribbean Coast of Panama and in the Gulf of Mexico: composition, structure and community ecology." *Marine Biology* 50: 97-107.
- Whaley, R. (1990). Causes of seagrass decline in the Tampa Bay area.

Zieman, J. C. (1976). "The ecological effects of physical damage from motor boats on turtle grass beds in Southern Florida." *Aquatic Botany* 2: 127-139.

Zimmermann, C. F., and Montgomery, J. R. (1984). "Effects of a decomposing drift algae mat on sediment pore water nutrient concentrations in a Florida seagrass bed." *Marine Ecology Progress Series* 19: 299-302.